



## **Modeling pesticides emissions for Grapevine LCA: adaptation of Pest-LCI model to viticulture**

**Christel, Renaud-Gentié ; Dijkman, Teunis Johannes; Bjørn, Anders; Birkved, Morten**

*Publication date:*  
2014

[Link back to DTU Orbit](#)

### *Citation (APA):*

Christel, R-G. (Author), Dijkman, T. J. (Author), Bjørn, A. (Author), & Birkved, M. (Author). (2014). Modeling pesticides emissions for Grapevine LCA: adaptation of Pest-LCI model to viticulture. Sound/Visual production (digital)

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# Modeling pesticide emissions for Grapevine Life Cycle Assessment: adaptation of PestLCI model to viticulture

**Christel Renaud-Gentié<sup>1,\*</sup>, Teunis J. Dijkman<sup>2</sup>, Anders Bjorn<sup>2</sup>, Morten Birkved<sup>2</sup>**

<sup>1</sup> UPSP GRAPPE, UMT VINITERA, SFR Quasav, Groupe ESA, Angers, France

<sup>2</sup> Technical University of Denmark, DTU Management Engineering, Lyngby, Denmark

**\*Corresponding author. E-mail: [c.renaud@groupe-esa.com](mailto:c.renaud@groupe-esa.com)**



Area under vines (2012) :



7.5Mha



3Mha

**Vine very sensitive to pests and diseases...**

**High pesticides active ingredients (PAI) consumption**

**Need to address the lack of viticulture specific pesticide emission LCI model,**

**We propose**

**A PestLCI 2.0 version tailored to appropriately account for viticulture specificities**

**USETox<sup>TM</sup> specific characterization factors**


**PestLCI 2.0/USETox<sup>TM</sup> combination on 2 cases**

**Comparison to 2 static approaches**

## Method: Pest-LCI 2.0 model


PestLCI - calculates the primary distribution of p.a.i.s between air, crop and soil  
- estimates p.a.i.s. fate to air, surface water, groundwater

(Birkved et al. 2006, Dijkman et al. 2012)





# Method: Addressing specificities of viticulture








# Method

## I distribution : drift, specific sprayers



Sprayers: drift curves introduced in PestLCI from Codis et al. 2011



Codis et al. 2011

Plus Tunnel sprayer (Ganzelmeier 2000)



Picture Codis et al 2011



Picture Dagnaud




# Method : I distribution, double cropping system

- Primary distribution process in PestLCI : wind drift ( $f_d$ ), pesticide deposition on soil ( $f_s$ ) pesticide deposition on leaves ( $f_l$ )

$$f_d + f_s + f_l = 1$$

- Addition of cover crop interception :  $f_l = f_{\text{vine}} + (f_{\text{covercrop}} * f_s * \% \text{covered})$
- 36  $f_l$  combinations



Vine Stage	density of cover crop canopy	% of soil surface covered by cover crop	$f_{\text{vine}}$	Fraction intercepted by cover crop (calculation)	$f_l$
0	none	0	0.1	0	0.10
III	average (50%)	100%	0.65	0.05	0.70




# New p.a.i.s in PestLCI 2.0 and USETox™ characterization factors



PestLCI models organic p.a.i.s fate based on physical and chemical properties of the substances

**29 p.a.i.s were added in PestLCI 2.0**



Molecular weight (g/mol)
Molar volume (g/mol)
Solubility in water (g/l)
Ref. Temp solubility (0C)
Vapour pressure (Pa)
Ref. Temp vapour pressure
pKa
Log Kow
Koc (l/kg)
Soil t½ (days)
Reference temp for pesticide biodegradation (oC)
Atmospheric OH rate (days) (cm3/molecules*sec)
Atmospheric OH rate ref temp (0C)
No sprayzone width (m)

Usetox™, best application practice to characterize toxic impacts of chemicals in LCA (Haushild et al. 2008), covers ~ 2500 chemicals for Fresh water ecotoxicity (FwEtox)

**18 p.a.i.s CF were calculated for FwEtox**





# Case study

2 real cases from regional typology of diversity of vineyard technical management: conventional , with different intensities of intervention,  
 Chenin Blanc, year 2011, different soils.

Functional Unit : 1ha

TMR 1  
 12.6 kg p.a.i.s  
 14 applications


TMR 3  
 4.17 kg p.a.i.s  
 11 applications

pesticide ai.s	Application rate	Crop type + development stage	Month of application	Application method	width treated
Aminotriazole	0,79	Grass I - all phases	april	sheltered boom	1,85
Aclonifen	0,31	Grass I - all phases	april	sheltered boom	1,85
Sulfur	5,89	*Vines II - h80% grass	may	tunnel sprayer	1,85
Folpel	0,74	*Vines II - h80% grass	may	tunnel sprayer	1,85
Fosétyl-Aluminium	1,47	*Vines II - h80% grass	may	tunnel sprayer	1,85
Fluopicolide	0,12	*Vines II - h80% grass	may	airblast sprayer	7,4
Fosétyl-Aluminium	1,75	*Vines II - h80% grass	may	airblast sprayer	7,4
Proquinazid Technique	0,05	*Vines II - h80% grass	may	airblast sprayer	7,4
Tétraconazole	0,03	*Vines III - a80% grass	june	airblast sprayer	7,4
<a href="#">Indoxacarbe</a>	0,04	*Vines III - a80% grass	june	airblast sprayer	7,4
<a href="#">copper oxychloride</a>	0,73	*Vines III - a80% grass	july	airblast sprayer	7,4
<a href="#">copper sulfate</a>	0,18	*Vines III - a80% grass	july	airblast sprayer	7,4
cymoxanil	0,12	*Vines III - a80% grass	july	airblast sprayer	7,4
Mancozèbe	0,40	*Vines III - a80% grass	july	airblast sprayer	7,4
Glyphosate	0,54	Grass I - all phases	march	sheltered boom	1,95
<a href="#">Amitrole</a>	0,92	Grass I - all phases	march	sheltered boom	1,95
ammonium thiocyanate	0,86	Grass I - all phases	march	sheltered boom	1,95
<a href="#">Flazasulfuron</a>	0,02	Grass I - all phases	march	sheltered boom	1,95
Glyphosate	0,09	Grass I - all phases	may	sheltered boom	1,95
<a href="#">Trifloxystrobin</a>	0,06	*Vines II - a50% grass	may	pneumatic sprayer side by side	7,8
<a href="#">Trifloxystrobin</a>	0,06	*Vines III - a50% grass	june	pneumatic sprayer side by side	7,8
<a href="#">Diméthomorph</a>	0,18	*Vines III - a50% grass	june	pneumatic sprayer side by side	7,8
<a href="#">Mancozèbe</a>	1,20	*Vines III - a50% grass	june	pneumatic sprayer side by side	7,8
<a href="#">Difénoconazole</a>	0,03	*Vines III - a50% grass	july	pneumatic sprayer side by side	7,8
Meptyldinocap	0,21	*Vines III - a50% grass	july	pneumatic sprayer side by side	7,8




# Results and discussion

## Emissions fractions



## Emissions quantities and FwEtox impacts




- results consistent with literature (Vazquez-Rowe et al. 2012)
- Need of sensitivity analysis on various parameters



# Results and discussion

Comparison with static approaches:

- 100% in agric. soil (Nemecek *et al.*, 2012)
- 75% soil, 25% air (Neto *et al.*, 2012)



Impacts	average impact
Impacts FwEcotox PestLCI 2.0 (PAF·m3·day)	484
impacts FwEcotox Ecoinvent (PAF·m3·day)	15 200
impacts FwEcotox Neto et al. (PAF·m3·day)	12 600



- ❖ PestLCI 2.0 was adapted without compromising the model framework
  - ❖ Inorganic p.a.i.s are not covered by PestLCI 2.0 > need for a specific model to permit organic farming assessment
  - ❖ Not included, consider for further improvement:
    - ❖ Airborne drift
    - ❖ p.a.i.s metabolites
    - ❖ Spraying conditions
    - ❖ application parameters
    - ❖ % stones in the soil
- difficult to model and/or lack of existing references
- More complicated inventory

- ✓ **Accounts for main viticulture specificities**
- ✓ Provides **emission and characterization factors** for main vineyards specific p.a.i.s
- ✓ Application to cases showed **high variations between p.a.i.s applications** due to p.a.i.s properties and application conditions
- ✓ Comparison to **static approaches** points out **huge differences of p.a.i.s emitted quantities and FwEtox** due to different system boundaries and phenomenon and a.i. properties accounting
- ✓ Can be used by **other vertical canopy crops**



# Aknowledgements



The authors thank


Interloire and Region pays de la Loire for funding,

The winegrowers for their time and data,

D. Boudiaf for her help in data collection,

F. Jourjon, R Siret, S Beauchet and E. Mehinagic for  
their core reading and advices





**Thank you for your attention**

**Questions, discussion**